

Andermatt

COMBI-PROTEC
INSECT BAIT ATTRACTANT

COMBI-PROTEC is a feed stimulating tool for fruit fly management that improves the efficiency and long-term effectiveness of insecticides in a tank mix.

Principal Functioning Agents: Plant Extracts **80.0%** | Sugars **20.0%**

This product is exempt from FIFRA regulation at 40 C. F. R. § 152.25 (d) "Foods".

Enhanced Efficacy

- Stimulates feeding which results in increasing oral insecticide uptake and accelerated mortality.
- Fast mortality reduces oviposition resulting in fewer rejects and better fruit quality.
- Minimizes the initial activity delay of insecticides.
- Compatibility with insecticides allows for easy rotation of actives.
- Less residue on flowers and fruit & more applications within seasonal limits.
- Fits full-coverage, bait-spray and low-volume programs
- Bait spray cuts water use up to 70% (100 → 30 GPA); less refilling for handheld and tractor-mounted sprayers.

Crop	Targeted Pest
Blackberry, Raspberry, Blueberry, Strawberry, Elderberry, Plums, Apricots, Peach, Nectarines, Grapes & other Susceptible Crops	Spotted Wing Drosophila & other Fruit Fly Species
Cherry	Spotted Wing Drosophila & Cherry Fruit Fly
Walnut	Walnut Husk Fly
Peach, Nectarines & other Susceptible Crops	Mediterranean Fruit Fly
Olive	Olive Fruit Fly
Hazelnut, Peach, Almonds & other Susceptible Crops	Brown Marmorated Stink Bug

Application, Timing, & Rates

General Mix in cold or warm water, up to 104 °F with agitation. Begin applications at first adult emergence and reapply every 5-10 days until harvest. Spray 24-48 hrs before high-risk events (rain, overhead irrigation).

- **Full Coverage** 1 lb / acre in a minimum of 10 gal water/acre for thorough canopy coverage. May be tank mixed with compatible fungicides, adjuvants, and foliar nutrients.
- **Bait Spray** 1-2 lb / acre plus a suitable insecticide, at ≥2.5 gal water per 1 lb product (≤30 GPA recommended). Medium-very coarse droplets: 300-500 µm for SWD, up to 4000 µm for other fruit flies. Only 3-10% foliage coverage needed. No tank mixes beyond the selected insecticide.
- **Aerial Coverage** 1-2 lb / acre plus a suitable insecticide, at ≥2.5 gal water per 1 lb product. Coarse droplet size must be maintained.

Follow the co-applied insecticide label (volume, rate) unless eligible for FIFRA §2 (ee) exemption. Confirm exemption with an extension specialist or CCA.



Restrictions

- Always tank-mix with a suitable insecticide. Never apply as a standalone product.
- Not compatible with stickers, extenders, adjuvants or additives that reduce droplet adhesion.
- Do not apply at the same time as, or shortly after, copper compounds.

Enhances Insecticide Performance | **Flexible Application Methods** | **Supports Lower Use Rates** | **Feeding Stimulant and Attractant** | **Compatible with Crop Programs** | **Suitable for Organic & Conventional Systems**

Andermatt USA Corporation
107 Gilbreth Parkway, Mullica Hill, NJ 08062
(302) 724-6888 · contact@andermattusa.com · www.andermattusa.com

©2024 dedetec Gmbh, Am Reichenbach 10, D 79249 Merzhausen (Germany) info@dedetec.de, ww.combi-protec.com. Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status. PIEN2024V240205